Accepted Manuscript




# *Ovar-DRB1* haplotypes *0404 and *0141 are associated with sheep growth and ewe lifetime prolificacy

Mehmet Ulas Cinar, Michelle R. Mousel, Lynn M. Herrmann-Hoesing, J. Bret Taylor, Stephen N. White

PII: S0378-1119(16)30798-3
DOI: doi:10.1016/j.gene.2016.10.004
Reference: GENE 41613

To appear in: *Gene*

Received date: 6 May 2016
Revised date: 15 August 2016
Accepted date: 1 October 2016

Please cite this article as: Cinar, Mehmet Ulas, Mousel, Michelle R., Herrmann-Hoesing, Lynn M., Taylor, J. Bret, White, Stephen N., *Ovar-DRB1* haplotypes *0404 and *0141 are associated with sheep growth and ewe lifetime prolificacy, *Gene* (2016), doi:10.1016/j.gene.2016.10.004

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

# *Ovar-DRB1* haplotypes *0404 and *0141 are associated with sheep growth and ewe lifetime prolificacy

**Mehmet Ulas Cinar[a,b], Michelle R. Mousel[c], Lynn M. Herrmann-Hoesing[a], J. Bret Taylor[d], Stephen N. White[a,c,e,*]**

a Department of Veterinary Microbiology and Pathology, Washington State University, Pullman, WA, USA

b Erciyes University, Faculty of Agriculture, Department of Animal Science, 38039 Kayseri, Republic of Turkey

c Animal Disease Research Unit, Agricultural Research Service, U.S. Department of Agriculture, Pullman, WA, USA

d Range Sheep Production Efficiency Research Unit, Agricultural Research Service, U.S. Department of Agriculture, Dubois, ID, USA

e Center for Reproductive Biology, Washington State University, Pullman, WA USA

**MUC**: ulas@vetmed.wsu.edu

**MRM**: michelle.mousel@ars.usda.gov

**LMHH**: lherrmann@vetmed.wsu.edu

**JBT**: bret.taylor@ars.usda.gov

**SNW**: stephen.white@ars.usda.gov

***Correspondence to:**

Dr. Stephen N. White

Animal Disease Research Unit, USDA-ARS, Pullman, WA, 99164, USA

E mail: stephen.white@ars.usda.gov

Phone: +1-509-335-7407

Fax: +1-509-335-8529

ACCEPTED MANUSCRIPT

ACCEPTED MANUSCRIPT

**Abstract**

Background: The major histocompatibility complex (MHC) is an organized cluster of tightly linked vertebrate genes with immunological and non-immunological functions. While the important MHC gene *DRB1* has been examined in regard to many sheep infectious disease traits, only one study, based on microsatellite markers, has previously examined *DRB1* and sheep production traits. Furthermore, to our knowledge no studies have examined *DRB1* relationship with lifetime ewe prolificacy traits. Therefore, we analyzed association between the presence of *DRB1* SNP haplotypes with internationally recognized standard names and production traits including growth and lifetime prolificacy in 370 Rambouillet, Columbia, and Polypay sheep.

Results: The *DRB1* *0404 haplotype was associated with increased weaning and mature weights, as well as average daily gain (Šidák $P < 0.05$; corrected for the number of haplotypes tested). Interestingly, the *0404 haplotype also showed a trend toward association with increased total number of lifetime lambs born (Šidák $P = 0.084$) and number of lambs born alive (Šidák $P = 0.084$). In contrast, the *DRB1* *0141 haplotype was associated with decreased mature weight (Šidák $P = 0.01$).

Conclusions: Since the *0404 haplotype was present in all three breeds, these results suggest there is at least one functional mutation in the region that influences growth and prolificacy traits that may be broadly present across several breeds. Furthermore, combined use of the similar *0404 and *0141 multi-marker haplotypes that nonetheless have opposing directions of production trait associations will enhance mutation discovery in this region. If undesirable alleles for underlying mutations can be identified, selective pressure against one or a small number of

ACCEPTED MANUSCRIPT

undesirable alleles may improve production with limited impact on MHC genetic diversity and infectious disease susceptibility.

Keywords

Major Histocompatibility Complex, MHC class II gene, Association analysis, Sheep, Production traits, Growth, Female Reproduction, Lifetime Prolificacy

Highlights

• *DRB1* *0404 haplotype was present in all breeds examined.

• *0404 haplotype was associated with increased growth and lifetime number of lambs born.

• Conversely, *0141 haplotype was associated with decreased growth.

• Contrast of *0404 and *0141 extended haplotypes will enhance mutation discovery.

## 1. Introduction

Sheep are a major domestic animal species for meat, milk, and wool products (Morris, 2009). The Major Histocompatibility Complex (MHC) of the sheep, designated as ovine Lymphocyte Antigen (OLA), harbors clusters of immunological genes involved in overall resistance/susceptibility of the animal to infectious diseases (Gao et al., 2010). Genetic loci in the MHC are organized to form distinct functional clusters as class I, II, and III, which show considerable conservation among mammal species, most likely due to their importance for host-pathogen interactions (Glynn, 1988). Class I and II MHC genes encode cell-surface glycoproteins (MHC molecules) that present small peptide antigens to T-cells, thereby controlling both cell- and antibody-mediated immune responses. The MHC class III region

ACCEPTED MANUSCRIPT

consists of genes sandwiched between the MHC class I and II regions in placental mammals and may play a role in the innate immune system as opposed to the adaptive immune system (Deakin et al., 2006). In addition, there are non-immunological genes in the MHC region, including components of the proteasome and iron metabolic pathways, among others (Janeway et al., 2001; Jiang et al., 2014).

Among sheep MHC class II genes, the *DRB1* and *DQB1* loci have been found to be highly polymorphic and this polymorphism is maintained by a form of balancing selection (Hughes and Yeager, 1998). In particular, exon 2 encodes the highly polymorphic antigen-binding site (Herrmann-Hoesing et al., 2008) where 106 *Ovar-DRB1* separate alleles have been identified (Ballingall et al., 1992; Fabb et al., 1993; Schwaiger et al., 1995; Kostia et al., 1998; Paterson, 1998; Nagaoka et al., 1999; Jugo and Vicario, 2000; Konnai et al., 2003a; Herrmann-Hoesing et al., 2008). Variation in the *Ovar-DRB1* gene may impact immune responses to pathogens, which may lead to variation in disease resistance. Numerous investigations have detected polymorphism of sheep MHC genes and described association with resistance or susceptibility to infectious diseases including studies on resistance to nematodiasis (Sayers et al., 2005; Stear et al., 2009; Valilou et al., 2015), bacterial diseases like footrot (Escayg et al., 1997) and viral diseases such as BLV-induced ovine lymphoma, small ruminant lentiviruses (SRLV: also called Maedi-Visna or ovine progressive pneumonia virus), and pulmonary adenocarcinoma viral diseases (Aida et al., 1995; Konnai et al., 2003b; Herrmann-Hoesing et al., 2008; Larruskain et al., 2010).

ACCEPTED MANUSCRIPT

Besides the role of the MHC in immune resistance, quantitative trait loci (QTL) have been identified in or near the MHC region associated with economically important traits such as growth and meat quality in pigs (Wei et al., 2011), semen traits and growth rate of young bulls (Grignola et al., 1995), increased protein yield and fat percentage in cow milk (Batra et al., 1989; Sharif et al., 1999), and milk yield, body weight, and maximum daily gain in sheep (Gutierrez-Gil et al., 2009; Hadjipavlou and Bishop, 2009; Raadsma et al., 2009). These QTL may be the result of pleiotropic effects of some MHC genes on physiologic processes, or they might result from linkage disequilibrium with other nearby genes. However, only one microsatellite-based study with 249 ewes and 381 lambs, found an association of *Ovar-DRB1* polymorphism with any growth and reproduction traits in sheep (Geldermann et al., 2006), and that study only used one year of records instead of ewe cumulative lifetime production. However, ewe cumulative lifetime production is one of the most economically important metrics, especially since replacement costs are a substantial fraction of total costs for ruminant livestock production systems. To the best of our knowledge, no study has evaluated an association of lifetime production traits with *Ovar-DRB1* in sheep. Therefore the aim of this study was to investigate association between *Ovar-DRB1* haplotypes and individual and lifetime production traits in Rambouillet, Columbia, and Polypay sheep.

## 2. Methods

### 2.1. Animals and phenotypes

We sampled 370 Rambouillet, Columbia, and Polypay ewes from the Range Sheep Production Efficiency Research station in Idaho as described previously (Herrmann-Hoesing et al., 2008). Briefly, approximately equal numbers were chosen by breed and age (3, 4, 5, and 6

ACCEPTED MANUSCRIPT

years in 2004) for a separate study on small ruminant lentivirus (Herrmann-Hoesing et al., 2008). In this analysis, the totals were: 129 Rambouillet, 117 Columbia, and 124 Polypay ewes. Phenotypes were recorded as individual traits and lifetime production traits (collected both before and after 2004), an accumulation of specific production traits. Individual ewe traits were defined as unadjusted birth weights (BRWT), unadjusted weights in the spring of the year born (JWT), unadjusted weaning weights (WNWT), weaning weight adjusted to a constant 120 days (WT120), unadjusted weights collected late summer in the year born (ASWT) and unadjusted weights twice a year during 3 and 4 years of age in spring and fall (SPWT3, FAWT3, SPWT4 and FAWT4). Average daily gain (ADG) was calculated as the difference between the unadjusted weight at weaning minus birth weight divided by an adjusted age of 120 days. For example, lambs younger than 120 days had an increased adjusted weaning weight and older lambs had a decreased adjusted weaning weight compared with unadjusted weaning weight. Fleece was shorn annually, and unwashed, greasy weight was recorded (LFLWT). Milk score (0 = none to 5 = very good) was a subjective assessment by trained staff on the amount of milk available shortly after birth in the final year of the ewe's productive life. Subjective udder (mammary gland) scores were collect each year with a 12 point scale in which 1 was removed from analysis because the ewe was not lactating, 2-3 were considered normal, and 4-12 were abnormal. Total number of lamb(s) born (LLB), number of lamb(s) born alive (LLBA), and number of lamb(s) weaned (LLW) were analyzed as cumulative lifetime counts. Cumulative ewe lifetime lamb birth weights were unadjusted (LLBW) and lifetime lamb weaning weights (LLWW) were adjusted to an average age of 120 days.

2.2. RNA isolation and genotyping of *Ovar-DRB1*

ACCEPTED MANUSCRIPT

Because distinguishing expressed genes from pseudogenes is an issue especially in the MHC (Kumanovics et al., 2003), we used RNA to generate genotypes only from expressed genes. Details of RNA isolation and genotyping of *Ovar-DRB1* were given in a previous study (Herrmann-Hoesing et al., 2008) that focused on different traits. Briefly, total RNA was extracted from 2 ml whole blood using PAXtube technology following manufacturer's directions (Qiagen, Hilden, Germany) and was stored at −80°C until needed. RNA quality was assessed by gel electrophoresis and 260/280 nm spectrophotometric analysis prior to RT-PCR, and RT-PCR positive and negative controls for each reaction set further confirmed input RNA quality of all samples. *Ovar-DRB1* allelic typing was conducted by cloning RT-PCR products spanning the peptide-binding site and sequencing, or by cDNA sequencing combined with use of electronic confirmation of known haplotypes (Stephens et al., 2001; Stephens and Donnelly, 2003; Herrmann-Hoesing et al., 2008), as previously described. The resulting haplotypes were matched to internationally accepted haplotype names (Herrmann-Hoesing et al., 2008; Ballingall et al., 2011), and these standardized names were used throughout the study.

### 2.3. Statistical analysis

Data were checked for normality before analyses using the UNIVARIATE procedure of SAS v9.4 (SAS Institute, Cary, NC). All individual and lifetime cumulative trait data were analyzed with a mixed model using the MIXED procedure of SAS v9.4. The model included fixed effects of breed, year of birth (corresponded to age in years for this study), and genotype, with a random effect of sire nested within breed. For discrete traits such as milk and udder scores, the GLIMMIX procedure of SAS v9.4 was used to examine association with *Ovar-DRB1* genotypes with the same fixed and random effects. A Šidák correction for multiple testing (Šidák, 1967)

ACCEPTED MANUSCRIPT

was applied on a haplotype basis for all haplotypes present in 30 or more individuals. The Šidák method for multiple testing correction makes the strong assumption of complete statistical independence, and it is therefore overly conservative in any set of analyses within which positive dependence occurs. For each statistical test, Šidák P-values were reported to account for multiple testing and and nominal *P* values were reported for completeness and to allow for additional reader analyses, if desired.

## 3. Results

Associations of *Ovar-DRB1* haplotypes with the individual and lifetime production traits in an Idaho ewe flock consisting of Rambouillet, Columbia, and Polypay breeds were investigated in 370 sheep. From sequencing of cDNA products, 31 DRB1 haplotypes were identified in 370 ewes. Total numbers of animals possessing at least one copy of each *DRB1* haplotype and overall breed-specific *DRB1* haplotype frequencies were presented in supplementary Table 1.

For individual ewe traits (excluding lamb production), the *DRB1* *0404 haplotype was significantly associated with JWT, WWT and ASWT after multiple testing (all Šidák $P < 0.05$; Table 1), and all three breeds carried this haplotype (supplementary Table 1). The *DRB1* *0404 haplotype also was nominally associated with WT120 ($P = 0.01$), ADG ($P = 0.01$), FAWT3 ($P = 0.04$) and showed a nominal trend toward association with FAWT4 ($P = 0.07$; Table 1). No additional significant associations with *DRB1* *0404 were found (supplementary Table 2). *DRB1* *0141 was nominally associated with WT120 ($P = 0.01$), WWT ($P = 0.01$), ADG ($P = 0.02$), ASWT ($P = 0.0081$), SPWT3 ($P = 0.0011$; Šidák $P = 0.01$), and FAWT3 ($P = 0.05$) (Table 2; supplementary Table 2). *DRB1* *0141 also showed a trend toward association with BWT

ACCEPTED MANUSCRIPT

($P = 0.09$), JWT ($P = 0.08$), and SPWT4 ($P = 0.07$) (supplementary Table 2). However, this haplotype was detected only in Columbia and Rambouillet ewes (supplementary Table 1). Nominally significant association was observed between several additional haplotypes and ewe individual traits that were not significant after Šidák correction for multiple testing (supplementary Table 2). None of the haplotypes was found to be associated with milk and udder traits (all nominal $P > 0.05$) and therefore were not shown in supplementary Table 2.

For ewe lifetime lamb production traits, *DRB1* *0404 was associated with LLBA and LLW (Table 1). *DRB1* *0141 was significantly association with LLB ($P = 0.03$) and a trend toward association with LLW ($P = 0.09$) and LLWW ($P = 0.1$; Table 2). Additional nominally significant results that were not significant after Šidák correction for multiple testing are shown in supplementary Table 3.

Additionally, thirty-two ewes in the population (8.6%) were homozygous for *DRB1* where 2.4%, 1.6%, and 1.1% of the population were homozygous for *DRB1* *02032, *DRB1* *1202, and *DRB1* *0206, respectively (Supplementary Table 4). In terms of breed-specific haplotypes, *DRB1* *GS1 was found only in Columbia ewes. *DRB1* *0803 and *DRB1* *J were only found in Rambouillet ewes. *DRB1* *1301 was found only in Polypay ewes (Supplementary Table 1). Summary statistics for individual and lifetime traits are in supplementary Table 5 are provided for context concerning the sheep population used.

## 4. Discussion

ACCEPTED MANUSCRIPT

In our study we identified several associations between *Ovar-DRB1* haplotypes and weight gain at different life stages as well as ewe lifetime number of lambs born and born alive (Tables 1 and 2; supplementary Tables 2 and 3). Lambs that grow rapidly and reach market weights at younger ages, require a shorter feeding period, have improved feed efficiency, and therefore, increase producer profits. Further, lamb survival is a complex trait influenced by direct genetic, maternal genetic, and environmental effects, and genetic selection to improve this trait is possible (Everett-Hincks et al., 2014).

In at least two previous experiments, Rambouillet, Columbia, and Polypay breeds were clearly clustered separately based on genome-wide analysis of 50,000 SNP (White et al., 2012; Zhang et al., 2013). This study identified breed differences in production relevant phenotypes. Specifically, we identified differences in birth weight, fleece weight, and lifetime number of lambs born (Tables 1 and 2). These findings are consistent with breed standards for a semi-prolific breed (Polypay), a wool breed (Rambouillet), and a multipurpose breed (Columbia) and with previous work (Bromley et al., 2000; Bromley et al., 2001). The breed differences in genome-wide genotypes and multiple phenotypic measures highlight genetic associations with the *0404 haplotype that was observed in all three breeds (supplementary Table 1).

However, there is very little literature regarding *Ovar-DRB1* association with production traits in sheep. Geldermann *et al.* (Geldermann et al., 2006) genotyped *OVA-DRB1* intron 2 microsatellite markers in 249 Merinoland ewes and found significant association with birth weight, weaning weight, and average daily gain. This is generally consistent with our findings, though growth trait associations reported here included more weights at older ages, in addition to

ACCEPTED MANUSCRIPT

the improved genetic markers from defined haplotypes. Major Histocompatibility Complex class II molecules are important cell-surface glycoproteins for the adaptive immune system because they present pathogen-derived peptides to CD4+ T lymphocytes (Duraes et al., 2013). While it is not immediately clear how MHC class II genes might be directly related with increase in body size, there are at least three possible explanations for indirect relationships of *DRB1* haplotype *0404 with growth and prolificacy traits shown here: 1) linkage disequilibrium with one or more other genes on ovine chromosome 20, 2) influence on susceptibility to one or more common pathogens with impact on production traits, and/or 3) less obvious mechanisms including downstream effects of inflammation.

The MHC is characterized by long, ancient haplotypes spanning many genes (Raymond et al., 2005). Thus, it is possible that the *DRB1* *0404 and *0141 haplotypes are in linkage disequilibrium with one or more functional mutations in or affecting other genes on ovine chromosome 20. The sheep QTL database (Hu et al., 2016) contains several significant QTLs associated with morphostructural phenotypes, average daily gain, body weight at different ages and maximum daily gain (Hadjipavlou and Bishop, 2009). In addition, consistent with our findings Geldermann *et al.* (Geldermann et al., 2006) found significant association between *OVA-DRB1* intron 2 variants and number of lambs born and number of lambs weaned in a single production year from Merinoland breed ewes. However, most of these studies used microsatellite markers that cannot be directly transferred across populations because of the high microsatellite mutation rate even within families. In contrast, the multi-marker haplotypes composed of evolutionarily stable SNP in named *DRB1* haplotypes *0404 and *0141 are readily identifiable in different populations. Thus, the specific identification of these named haplotypes will facilitate

ACCEPTED MANUSCRIPT

validation of our findings across populations, which is important for adoption in breeding programs and for mutation discovery (White and Knowles, 2013). Phylogenetic analysis of 31 *Ovar-DRB1* haplotypes revealed three main branches called A, B and C with *0404 and *0141 haplotypes found in branch C (Herrmann-Hoesing et al., 2008). This makes the opposing directions of association (Tables 1 and 2) between these similar *DRB1* haplotypes more useful, because comparing similar extended haplotypes may help greatly reduce the search region for potential causal mutations. Additionally the region of ovine chromosome 20 near *Ovar-DRB1* harbors genes related with reproduction traits such as glutathione S-transferase 1 and 4 (*GSTA1* and *GSTA4*) which were associated with oocyte competence (Salhab et al., 2011) and granulosa cell development (Fayad et al., 2004), as well as genes related with growth traits like tenascin XB (*TNXB*) (Ajayi et al., 2014).

A second hypothesis to explain the observed associations could involve improved control of one or more common pathogens with consequent improved average growth. Indeed, the associations we observed with lifetime traits may suggest improved overall health for sheep bearing the *0404 haplotype. There are also several examples of data from the literature that may support this hypothesis. The *0404 haplotype was found only in healthy individuals but not in sheep with small ruminant lentiviruses (SRLV) or ovine pulmonary adenocarcinoma virus-related disease (Larruskain et al., 2010). Furthermore, the *0404 haplotype contains amino acid N74 that was previously associated with improved control of SRLV (reduced SRLV proviral concentration) (Herrmann-Hoesing et al., 2008). Both SRLV and pulmonary adenocarcinoma virus can affect growth (Arsenault et al., 2003; Bennett, 2003; Bennett and Ijpelaar, 2005). Association of *DRB1* variants with resistance to intestinal nematode infection were shown in

ACCEPTED MANUSCRIPT

several sheep breeds (Sayers et al., 2005; Stear et al., 2009; Shen et al., 2014; Valilou et al., 2015). Generally, increase in faecal egg count of nematodes is associated with decreased sheep performance in terms of body weight, wool production, and milk yield (Mavrot et al., 2015). For example, a population of Scottish Blackface sheep had QTLs in the vicinity of the *Ovar-DRB1* locus associated with parasitic resistance (Davies et al., 2006), daily gain, and body weight (Hadjipavlou and Bishop, 2009). These and other examples suggest that *DRB1* may be indirectly related to sheep production through improved pathogen control. If so, the six amino acid residues that differ between *0404 and *0141 may be implicated in differential presentation of important pathogen epitopes.

A third possibility might include downstream effects of inflammatory conditions, which can result in significant upregulation of MHC II molecules (Nagaraju, 2001). Skeletal muscles can be injured by their own contractions during everyday activities (Marino et al., 2011). . The range ewes we evaluated are required to walk up and down mountains for forage and water, and the exertion may contribute to muscle inflammation and help explain the association of *Ovar-DRB1* variants with growth traits in our ewe population.

The MHC plays a critical role in the immune recognition of parasites and pathogens. There are possible associations of MHC loci with a number of quantitative traits linked to the fitness and behavior of individuals in natural populations (Piertney and Oliver, 2006). A genetic selection program based on MHC or nearby genes may cause undesirable effects on diversity maintained over long periods of balancing selection. However, Apanius' heterozygote advantage and frequency dependence theories (Apanius et al., 1997) suggest that eliminating the most

ACCEPTED MANUSCRIPT

susceptible alleles from the global population, but keeping other alleles, could be effective for livestock populations with minimal impact on overall MHC diversity. Indeed, (Maillard et al., 2003) demonstrated a practical example of such selection and defeated a population-limiting infectious disease crisis in cattle. If underlying mutations affecting growth and reproductive performance can be found on ovine chromosome 20, elimination or reduction of some undesirable performance alleles could be implemented with limited impact on flock level MHC genetic diversity and infectious disease resistance.

## 5. Conclusion

In summary, our results define associations between *0404 and *0141 *Ovar-DRB1* haplotypes and growth and prolificacy traits across 3 different sheep breeds. Specifically, the *DRB1* *0404 haplotype was associated with increased weaning and mature weights, as well as average daily gain, while the *DRB1* *0141 haplotype was associated with decreased mature weight. Interestingly, the *0404 haplotype also showed a trend toward association with increased total number of lifetime lambs born and number of lambs born alive. These observed associations suggest the existence of one or more functional mutations on the *DRB1* *0404 and *0141 haplotypes which affect prolificacy and growth performance. Further studies will be required in order to understand sheep MHC genetics and identify functional mutations on ovine chromosome 20 involved in reproduction, prolificacy, growth, and other production traits.

## Conflict of interest

The authors declare they have no conflicts of interest.

ACCEPTED MANUSCRIPT

**Acknowledgements**

We thank Nic Durfee, Codie Durfee, Margaret Highland, Ralph Horn, James Allison, Megan Blauert, Lori Fuller, James Reynolds and ADRU staff for technical assistance. We also wish to thank current and former RSPER employees Tom Kellom and Natalie Pierce for data collection, curation, and management assistance, and Shiquan Wang, Mark Williams, Boyd Leonard, John Mackin, Ella Ybarlucea, Harley Carpenter, Joel Billman, Brad Eddins, Lyn Mortensen, and Jack Hensley for animal management, husbandry, and sample-collection assistance. We also thank former University of Idaho employee M. Quinn Jacobson for animal management, husbandry, and sample-collection assistance. This work was supported by USDA-ARS Cwa. 2090-32000-031-00D. The funder had no role in study design; in collection, analysis or interpretation of data; in writing the report; or in the decision to submit the article for publication.

**References**

Aida, Y., Niimi, M., Asahina, M., Okada, K., Nakai, Y. and Ogimoto, K., 1995. Identification of a new bovine MHC class II DRB allele by nucleotide sequencing and an analysis of phylogenetic relationships. Biochem Biophys Res Commun 209, 981-988.

Ajayi, O.O., Adefenwa, M.A., Agaviezor, B.O., Ikeobi, C.O.N., Wheto, M., Okpeku, M., Amusan, S.A., Yakubu, A., De Donato, M., Peters, S.O. and Imumorin, I.G., 2014. A novel TaqI polymorphism in the coding region of the ovine TNXB gene in the MHC Class III region: Morphostructural and physiological influences. Biochem Genet 52, 1-14.

Apanius, V., Penn, D., Slev, P.R., Ruff, L.R. and Potts, W.K., 1997. The nature of selection on the major histocompatibility complex. Crit Rev Immunol 17, 179-224.

Arsenault, J., Dubreuil, P., Girard, C., Simard, C. and Belanger, D., 2003. Maedi-visna impact on productivity in Quebec sheep flocks (Canada). Prev Vet Med 59, 125-137.

Ballingall, K.T., Herrmann-Hoesing, L., Robinson, J., Marsh, S.G. and Stear, M.J., 2011. A single nomenclature and associated database for alleles at the major histocompatibility complex class II DRB1 locus of sheep. Tissue Antigens 77, 546-553.

Ballingall, K.T., Wright, H., Redmond, J., Dutia, B.M., Hopkins, J., Lang, J., Deverson, E.V., Howard, J.C., Puri, N. and Haig, D., 1992. Expression and characterization of ovine major histocompatibility complex class II (OLA-DR) genes. Anim Genet 23, 347-359.

ACCEPTED MANUSCRIPT

Batra, T.R., Lee, A.J., Gavora, J.S. and Stear, M.J., 1989. Class I alleles of the bovine major histocompatibility system and their association with economic traits. J Dairy Sci 72, 2115-2124.

Bennett, R., 2003. The 'direct costs' of livestock disease: The development of a system of models for the analysis of 30 endemic livestock diseases in Great Britain. J Agricult Econ 54, 55-71.

Bennett, R. and Ijpelaar, J., 2005. Updated estimates of the costs associated with thirty four endemic livestock diseases in Great Britain: A note. J Agricult Econ 56, 135-144.

Bromley, C.M., Snowder, G.D. and Van Vleck, L.D., 2000. Genetic parameters among weight, prolificacy, and wool traits of Columbia, Polypay, Rambouillet, and Targhee sheep. J Anim Sci 78, 846-858.

Bromley, C.M., Van Vleck, L.D. and Snowder, G.D., 2001. Genetic correlations for litter weight weaned with growth, prolificacy, and wool traits in Columbia, Polypay, Rambouillet, and Targhee sheep. J Anim Sci 79, 339-346.

Davies, G., Stear, M.J., Benothman, M., Abuagob, O., Kerr, A., Mitchell, S. and Bishop, S.C., 2006. Quantitative trait loci associated with parasitic infection in Scottish blackface sheep. Heredity 96, 252-258.

Deakin, J.E., Papenfuss, A.T., Belov, K., Cross, J.G.R., Coggill, P., Palmer, S., Sims, S., Speed, T.P., Beck, S. and Graves, J.A.M., 2006. Evolution and comparative analysis of the MHC Class III inflammatory region. BMC Genomics 7, 281.

Duraes, F.V., Thelemann, C., Sarter, K., Acha-Orbea, H., Hugues, S. and Reith, W., 2013. Role of major histocompatibility complex class II expression by non-hematopoietic cells in autoimmune and inflammatory disorders: facts and fiction. Tissue Antigens 82, 1-15.

Escayg, A.P., Hickford, J.G. and Bullock, D.W., 1997. Association between alleles of the ovine major histocompatibility complex and resistance to footrot. Res Vet Sci 63, 283-287.

Everett-Hincks, J.M., Mathias-Davis, H.C., Greer, G.J., Auvray, B.A. and Dodds, K.G., 2014. Genetic parameters for lamb birth weight, survival and death risk traits. J Anim Sci 92, 2885-2895.

Fabb, S.A., Maddox, J.F., Gogolin-Ewens, K.J., Baker, L., Wu, M.J. and Brandon, M.R., 1993. Isolation, characterization and evolution of ovine major histocompatibility complex class II DRA and DQA genes. Anim Genet 24, 249-255.

Fayad, T., Levesque, V., Sirois, J., Silversides, D.W. and Lussier, J.G., 2004. Gene expression profiling of differentially expressed genes in granulosa cells of bovine dominant follicles using suppression subtractive hybridization. Biol Reprod 70, 523-533.

Gao, J., Liu, K., Liu, H., Blair, H.T., Li, G., Chen, C., Tan, P. and Ma, R.Z., 2010. A complete DNA sequence map of the ovine major histocompatibility complex. BMC Genomics 11, 466.

Geldermann, H., Mir, M.R., Kuss, A.W. and Bartenschlager, H., 2006. OLA-DRB1 microsatellite variants are associated with ovine growth and reproduction traits. Genet Sel Evol 38, 431-444.

Glynn, L.E., 1988. Natural history of the major histocompatibility complex. Cell Biochem Funct 6, 222.

Grignola, F., Mao, I.L., Mejdell, C., Lie, O. and Solbu, H., 1995. Association of alleles for the class I bovine lymphocyte antigens with conformation, semen traits, and growth rate of young bulls. J Dairy Sci 78, 908-913.

ACCEPTED MANUSCRIPT

Gutierrez-Gil, B., El-Zarei, M.F., Alvarez, L., Bayon, Y., de la Fuente, L.F., San Primitivo, F. and Arranz, J.J., 2009. Quantitative trait loci underlying milk production traits in sheep. Anim Genet 40, 423-434.

Hadjipavlou, G. and Bishop, S.C., 2009. Age-dependent quantitative trait loci affecting growth traits in Scottish Blackface sheep. Anim Genet 40, 165-175.

Herrmann-Hoesing, L.M., White, S.N., Mousel, M.R., Lewis, G.S. and Knowles, D.P., 2008. Ovine progressive pneumonia provirus levels associate with breed and Ovar-DRB1. Immunogenetics 60, 749-758.

Hu, Z.-L., Park, C.A. and Reecy, J.M., 2016. Developmental progress and current status of the Animal QTLdb. Nucleic Acids Res 44, D827-D833.

Hughes, A.L. and Yeager, M., 1998. Natural selection and the evolutionary history of major histocompatibility complex loci. Front Biosci 3, d509-516.

Janeway, C.A., Travers, P., Walport, M. and Shlomchik, M.J., 2001. Immunobiology: the immune system in health and disease, 5th ed. Garland, New York.

Jiang, Y., Xie, M., Chen, W., Talbot, R., Maddox, J.F., Faraut, T., Wu, C., Muzny, D.M., Li, Y., Zhang, W., Stanton, J.A., Brauning, R., Barris, W.C., Hourlier, T., Aken, B.L., Searle, S.M., Adelson, D.L., Bian, C., Cam, G.R., Chen, Y., Cheng, S., DeSilva, U., Dixen, K., Dong, Y., Fan, G., Franklin, I.R., Fu, S., Fuentes-Utrilla, P., Guan, R., Highland, M.A., Holder, M.E., Huang, G., Ingham, A.B., Jhangiani, S.N., Kalra, D., Kovar, C.L., Lee, S.L., Liu, W., Liu, X., Lu, C., Lv, T., Mathew, T., McWilliam, S., Menzies, M., Pan, S., Robelin, D., Servin, B., Townley, D., Wang, W., Wei, B., White, S.N., Yang, X., Ye, C., Yue, Y., Zeng, P., Zhou, Q., Hansen, J.B., Kristiansen, K., Gibbs, R.A., Flicek, P., Warkup, C.C., Jones, H.E., Oddy, V.H., Nicholas, F.W., McEwan, J.C., Kijas, J.W., Wang, J., Worley, K.C., Archibald, A.L., Cockett, N., Xu, X., Wang, W. and Dalrymple, B.P., 2014. The sheep genome illuminates biology of the rumen and lipid metabolism. Science 344, 1168-1173.

Jugo, B.M. and Vicario, A., 2000. Single-strand conformational polymorphism and sequence polymorphism of Mhc-DRB in Latxa and Karrantzar sheep: implications for Caprinae phylogeny. Immunogenetics 51, 887-897.

Konnai, S., Nagaoka, Y., Takesima, S., Onuma, M. and Aida, Y., 2003a. Technical note: DNA typing for ovine MHC DRB1 using polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP). J Dairy Sci 86, 3362-3365.

Konnai, S., Takeshima, S.N., Tajima, S., Yin, S.A., Okada, K., Onuma, M. and Aida, Y., 2003b. The influence of ovine MHC class II DRB1 alleles on immune response in bovine leukemia virus infection. Microbiol Immunol 47, 223-232.

Kostia, S., Kantanen, J., Kolkkala, M. and Varvio, S.L., 1998. Applicability of SSCP analysis for MHC genotyping: fingerprinting of Ovar-DRB1 exon 2 alleles from Finnish and Russian breeds. Anim Genet 29, 453-455.

Kumanovics, A., Takada, T. and Lindahl, K.F., 2003. Genomic organization of the mammalian MHC. Annu Rev Immunol 21, 629-657.

Larruskain, A., Minguijon, E., Garcia-Etxebarria, K., Moreno, B., Arostegui, I., Juste, R.A. and Jugo, B.M., 2010. MHC class II DRB1 gene polymorphism in the pathogenesis of Maedi-Visna and pulmonary adenocarcinoma viral diseases in sheep. Immunogenetics 62, 75-83.

ACCEPTED MANUSCRIPT

Maillard, J.C., Berthier, D., Chantal, I., Thevenon, S., Sidibe, I., Stachurski, F., Belemsaga, D., Razafindraibe, H. and Elsen, J.M., 2003. Selection assisted by a BoLA-DR/DQ haplotype against susceptibility to bovine dermatophilosis. Genet Sel Evol 35, S193-S200.

Marino, M., Scuderi, F., Provenzano, C. and Bartoccioni, E., 2011. Skeletal muscle cells: from local inflammatory response to active immunity. Gene Ther 18, 109-116.

Mavrot, F., Hertzberg, H. and Torgerson, P., 2015. Effect of gastro-intestinal nematode infection on sheep performance: a systematic review and meta-analysis. Parasit Vectors 8, 557.

Morris, S.T., 2009. Economics of sheep production. Small Rumin Res 86, 59-62.

Nagaoka, Y., Kabeya, H., Onuma, M., Kasai, N., Okada, K. and Aida, Y., 1999. Ovine MHC class II DRB1 alleles associated with resistance or susceptibility to development of bovine leukemia virus-induced ovine lymphoma. Cancer Res 59, 975-981.

Nagaraju, K., 2001. Immunological capabilities of skeletal muscle cells. Acta Physiol Scand 171, 215-223.

Paterson, S., 1998. Evidence for balancing selection at the major histocompatibility complex in a free-living ruminant. J Hered 89, 289-294.

Piertney, S.B. and Oliver, M.K., 2006. The evolutionary ecology of the major histocompatibility complex. Heredity 96, 7-21.

Raadsma, H.W., Jonas, E., McGill, D., Hobbs, M., Lam, M.K. and Thomson, P.C., 2009. Mapping quantitative trait loci (QTL) in sheep. II. Meta-assembly and identification of novel QTL for milk production traits in sheep. Genet Sel Evol 41, 45.

Raymond, C.K., Kas, A., Paddock, M., Qiu, R., Zhou, Y., Subramanian, S., Chang, J., Palmieri, A., Haugen, E., Kaul, R. and Olson, M.V., 2005. Ancient haplotypes of the HLA Class II region. Genome Res 15, 1250-1257.

Salhab, M., Tosca, L., Cabau, C., Papillier, P., Perreau, C., Dupont, J., Merrnillod, P. and Uzbekova, S., 2011. Kinetics of gene expression and signaling in bovine cumulus cells throughout IVM in different mediums in relation to oocyte developmental competence, cumulus apoptosis and progesterone secretion. Theriogenology 75, 90-104.

Sayers, G., Good, B., Hanrahan, J.P., Ryan, M., Angles, J.M. and Sweeney, T., 2005. Major histocompatibility complex DRB1 gene: its role in nematode resistance in Suffolk and Texel sheep breeds. Parasitology 131, 403-409.

Schwaiger, F.W., Gostomski, D., Stear, M.J., Duncan, J.L., McKellar, Q.A., Epplen, J.T. and Buitkamp, J., 1995. An ovine major histocompatibility complex DRB1 allele is associated with low faecal egg counts following natural, predominantly Ostertagia circumcincta infection. Int J Parasitol 25, 815-822.

Sharif, S., Mallard, B.A., Wilkie, B.N., Sargeant, J.M., Scott, H.M., Dekkers, J.C. and Leslie, K.E., 1999. Associations of the bovine major histocompatibility complex DRB3 (BoLA-DRB3) with production traits in Canadian dairy cattle. Anim Genet 30, 157-160.

Shen, H., Han, G., Jia, B., Jiang, S. and Du, Y., 2014. MHC-DRB1/DQB1 Gene Polymorphism and Its Association with Resistance/Susceptibility to Cystic Echinococcosis in Chinese Merino Sheep. J Parasitol Res 2014, 272601.

Šidák, Z., 1967. Rectangular Confidence Regions for the Means of Multivariate Normal Distributions. J Am Stat Assoc 62, 626-633.

Stear, M.J., Boag, B., Cattadori, I. and Murphy, L., 2009. Genetic variation in resistance to mixed, predominantly Teladorsagia circumcincta nematode infections of sheep: from heritabilities to gene identification. Parasite Immunol 31, 274-282.

ACCEPTED MANUSCRIPT

Stephens, M. and Donnelly, P., 2003. A comparison of bayesian methods for haplotype reconstruction from population genotype data. Am J Hum Genet 73, 1162-1169.

Stephens, M., Smith, N.J. and Donnelly, P., 2001. A new statistical method for haplotype reconstruction from population data. Am J Hum Genet 68, 978-989.

Valilou, R.H., Rafat, S.A., Notter, D.R., Shojda, D., Moghaddam, G. and Nematollahi, A., 2015. Fecal egg counts for gastrointestinal nematodes are associated with a polymorphism in the MHC-DRB1 gene in the Iranian Ghezel sheep breed. Front Genet 6, 105.

Wei, W.H., Skinner, T.M., Anderson, J.A., Southwood, O.I., Plastow, G., Archibald, A.L. and Haley, C.S., 2011. Mapping QTL in the porcine MHC region affecting fatness and growth traits in a Meishan/Large White composite population. Anim Genet 42, 83-85.

White, S.N. and Knowles, D.P., 2013. Expanding possibilities for intervention against small ruminant lentiviruses through genetic marker-assisted selective breeding. Viruses 5, 1466-1499.

White, S.N., Mousel, M.R., Herrmann-Hoesing, L.M., Reynolds, J.O., Leymaster, K.A., Neibergs, H.L., Lewis, G.S. and Knowles, D.P., 2012. Genome-wide association identifies multiple genomic regions associated with susceptibility to and control of ovine lentivirus. PLoS One 7, e47829.

Zhang, L., Mousel, M.R., Wu, X., Michal, J.J., Zhou, X. and Ding, B., 2013. Genome-wide genetic diversity and differentially selected regions among Suffolk, Rambouillet, Columbia, Polypay, and Targhee sheep. PLoS One 8, e65942.

ACCEPTED MANUSCRIPT

ACCEPTED MANUSCRIPT

Abbreviations

MHC, major histocompatibility complex; DRB1, DR beta 1; OLA, ovine lymphocyte antigen; DQB1, DQ beta 1; DNA, Deoxyribonucleic acid; BLV, Bovine leukemia virus; SRLV, Small ruminant lentiviruses; QTL, quantitative trait loci; cDNA, complementary DNA; RT-PCR, Reverse transcription polymerase chain reaction; SNP, single nucleotide polymorphism; GSTA1, glutathione S-transferase 1; GSTA4, glutathione S-transferase 4; °C, centigrade.

BRWT, birth weight; JWT, weight in the spring of the year born; WNWT, weaning weight; WT120, weaning weight adjusted to a constant 120 days; ASWT, weight collected late summer in the year born; SPWT3, weight of 3 years of age in spring; SPWT4, weight of 4 years of age in spring; FAWT3, weight of 3 years of age in fall; FAWT4, weight of 4 years of age in fall; ADG, average daily gain; LLB, total number of lambs born; LLBA, number of lambs born alive; LLW, number of lambs weaned; LLBW, cumulative ewe lifetime lamb birth weights; LLWW, lifetime lamb weaning weights; LFLWT, lifetime greasy fleece weight.

ACCEPTED MANUSCRIPT

ACCEPTED MANUSCRIPT

**Table 1.** Association of *Ovar-DRB1* *0404 haplotype with individual and lifetime ewe traits.

| Trait | Presence of haplotype | Absence of haplotype | Nominal *P*-value | Šidák *P*-value |
|---|---|---|---|---|
| **JWT** [a] | 30.78 | 28.84 | 0.0031 | 0.040 |
| **WWT** [a] | 40.61 | 37.99 | 0.0003 | 0.004 |
| **ASWT** [a] | 46.10 | 43.37 | 0.0003 | 0.004 |
| **WT120** [a] | 40.27 | 38.46 | 0.014 | 0.14 |
| **ADG** [a] | 0.29 | 0.27 | 0.013 | 0.14 |
| **FAWT3** [a] | 82.21 | 79.48 | 0.047 | 0.47 |
| **FAWT4** [a] | 83.20 | 77.14 | 0.079 | 0.68 |
| **LLB** [b] | 12.57 | 11.52 | 0.0097 | 0.14 |
| **LLBA** [b] | 10.51 | 11.47 | 0.01 | 0.14 |

JWT: June weight after birth, WWT: Weaning weight, ASWT: August-September weight in the year born, WT120: Adjusted weight to a constant 120 days, ADG: Average daily gain, FAWT3: Weight in September of 3 years old, FAWT4: Weight in September of 4 years old. LLB: Lifetime number of lambs born; LLBA: Lifetime number of lambs born alive.

[a]Weight phenotype in kg.

[b]Lamb count phenotype.

ACCEPTED MANUSCRIPT

ACCEPTED MANUSCRIPT

**Table 2.** Association of *Ovar-DRB1* *0141 haplotype with individual and lifetime ewe traits.

| Trait | Presence of haplotype | Absence of haplotype | Nominal *P*-value | Šidák *P*-value |
|---|---|---|---|---|
| **SPWT3** [a] | 68.70 | 75.94 | 0.0011 | 0.01 |
| **BWT** [a] | 4.63 | 4.90 | 0.090 | 0.77 |
| **JWT** [a] | 27.53 | 29.08 | 0.087 | 0.73 |
| **WT120** [a] | 36.30 | 38.73 | 0.018 | 0.14 |
| **WWT** [a] | 35.85 | 38.34 | 0.014 | 0.14 |
| **ASWT** [a] | 40.95 | 43.75 | 0.0081 | 0.12 |
| **ADG** [a] | 0.26 | 0.28 | 0.020 | 0.27 |
| **FAWT3** [a] | 73.60 | 77.92 | 0.058 | 0.55 |
| **SPWT4** [a] | 68.70 | 75.94 | 0.071 | 0.68 |
| **LLB** [b] | 11.63 | 11.37 | 0.030 | 0.32 |
| **LLW** [b] | 10.73 | 10.60 | 0.094 | 0.77 |
| **LLWW** [a] | 8.75 | 8.63 | 0.10 | 0.81 |

SPWT3: Weight in April/May of 3 years old, BWT: Birth weight; JWT: June weight after birth, WT120: Adjusted weight to a constant 120 days, WWT: Weaning weight, ASWT: August-September weight in the year born, ADG: Average daily gain, FAWT3: Weight in September of 3 years old, FAWT4: Weight in September of 4 years old, LLB: Lifetime number of lambs born; LLW: Lifetime number of lambs weaned; LLWW: Total weight of lambs weaned over a lifetime.

[a]Weight phenotype in kg.

[b]Lamb count phenotype.